

Graph Editing to a Fixed Target^{*}

Petr A. Golovach¹, Daniël Paulusma², and Iain Stewart²

¹ Department of Informatics, Bergen University,
PB 7803, 5020 Bergen, Norway
`petr.golovach@ii.uib.no`

² School of Engineering and Computing Sciences, Durham University,
Science Laboratories, South Road, Durham DH1 3LE, United Kingdom
`{daniel.paulusma,i.a.stewart}@durham.ac.uk`

Abstract. For a fixed graph H , the H -MINOR EDIT problem takes as input a graph G and an integer k and asks whether G can be modified into H by a total of at most k edge contractions, edge deletions and vertex deletions. Replacing edge contractions by vertex dissolutions yields the H -TOPOLOGICAL MINOR EDIT problem. For each problem we show polynomial-time solvable and NP-complete cases depending on the choice of H . Moreover, when G is AT-free, chordal or planar, we show that H -MINOR EDIT is polynomial-time solvable for all graphs H .

1 Introduction

Graph editing problems are well studied both within algorithmic and structural graph theory and beyond (e.g. [1, 4, 22, 23]), as they capture numerous graph-theoretic problems with a variety of applications. A graph editing problem takes as input a graph G and an integer k , and the question is whether G can be modified into a graph that belongs to some prescribed graph class \mathcal{H} by using at most k operations of one or more specified types. So far, the most common graph operations that have been considered are vertex deletions, edge deletions and edge additions. Well-known problems obtained in this way are FEEDBACK VERTEX SET, ODD CYCLE TRANSVERSAL, MINIMUM FILL-IN, and CLUSTER EDITING. Recently, several papers [9, 10, 15–17] appeared that consider the setting in which the (only) permitted type of operation is that of an *edge contraction*. This operation removes the vertices u and v of the edge uv from the graph and replaces them by a new vertex that is made adjacent to precisely those remaining vertices to which u or v was previously adjacent. So far, the situation in which we allow edge contractions *together with* one or more additional types of graph operations has not been studied. This is the main setting that we consider in our paper.

A natural starting approach is to consider families of graphs \mathcal{H} of cardinality 1. For such families, straightforward polynomial-time algorithms exist if the set of permitted operations may only include edge additions, edge deletions and vertex deletions. However, we show that this is no longer necessarily true in

^{*} Research supported by EPSRC (EP/G043434/1) and ERC (267959).

our case, in which we allow both edge contractions and vertex deletions to be applied.

It so happens that setting $\mathcal{H} = \{H\}$ for some graph H , called the *target* graph from now on, yields graph editing problems that are closely related to problems that ask whether a given graph H appears as a “pattern” within another given graph G so that G can be transformed to H via a sequence of operations *without* setting a bound k on the number of operations allowed. These ‘unbounded’ problems are ubiquitous in computer science, and below we shortly survey a number of known results on them; those results that we will use in our proofs are stated as lemmas.

We start with some additional terminology. A *vertex dissolution* is the removal of a vertex v with exactly two neighbors u and w , which may not be adjacent to each other, followed by the inclusion of the edge uw . If we can obtain a graph H from a graph G by a sequence that on top of vertex deletions and edge deletions may contain operations of one additional type, namely edge contractions or vertex dissolutions, then G contains H as a *minor* or *topological minor*, respectively. For a *fixed* graph H , that is, H is not part of the input, this leads to the decision problems H -MINOR and H -TOPOLOGICAL MINOR, respectively. Grohe, Kawarabayashi, Marx, and Wollan [13] showed that H -TOPOLOGICAL MINOR can be solved in cubic time for all graphs H , whereas Robertson and Seymour [25] proved the following seminal result.

Lemma 1 ([25]). *H -MINOR can be solved in cubic time for all graphs H .*

We say that a containment relation is *induced* if edge deletions are excluded from the permitted graph operations. In the case of minors and topological minors, this leads to the corresponding notions of being an *induced minor* and *induced topological minor*, respectively, with corresponding decision problems H -INDUCED MINOR and H -INDUCED TOPOLOGICAL MINOR. In contrast to their non-induced counterparts, the complexity classifications of these two problems have not yet been settled. In fact, the complexity status of H -INDUCED MINOR when H is restricted to be a tree has been open since it was posed at the AMS-IMS-SIAM Joint Summer Research Conference on Graph Minors in 1991. Up until now, only forests on at most seven vertices have been classified [8] (with one forest still outstanding), and no NP-complete cases of forests H are known. The smallest known NP-complete case is the graph H^* on 68 vertices displayed in Figure 1; this result is due to Fellows, Kratochvíl, Middendorf and Pfeiffer [7].

Lemma 2 ([7]). *H^* -INDUCED MINOR is NP-complete.*

Lévêque, Lin, Maffray, and Trotignon [20] gave both polynomial-time solvable and NP-complete cases for H -INDUCED TOPOLOGICAL MINOR. In particular they showed the following result, where we denote the complete graph on n vertices by K_n .

Lemma 3 ([20]). *K_5 -INDUCED TOPOLOGICAL MINOR is NP-complete.*




Fig. 1. The smallest graph H^* for which H -INDUCED MINOR is NP-complete [7].

The complexity of H -INDUCED TOPOLOGICAL MINOR is still open when H is a complete graph on 4 vertices. L ev eque, Maffray, and Trotignon [21] gave a polynomial-time algorithm for recognizing graphs that neither contain K_4 as an induced topological minor nor a wheel as an induced subgraph. However, they explain that a stronger decomposition theorem (avoiding specific cutsets) is required to resolve the complexity status of K_4 -INDUCED TOPOLOGICAL MINOR affirmatively.

Before we present our results, we first introduce some extra terminology. Let G be a graph and H a minor of G . Then a sequence of minor operations that modifies G into H is called an H -minor sequence or just a minor sequence of G if no confusion is possible. The *length* of an H -minor sequence is the number of its operations. An H -minor sequence is *minimum* if it has minimum length over all H -minor sequences of G . For a fixed graph H , the H -MINOR EDIT problem is that of testing whether a given graph G has an H -minor sequence of length at most k for some given integer k . Also, for the other containment relations we define such a sequence and corresponding decision problem.

Because any vertex deletion, vertex dissolution and edge contraction reduces a graph by exactly one vertex, any H -induced minor sequence and any H -topological induced minor sequence of a graph G has the same length for any graph H , namely $|V_G| - |V_H|$. Hence, H -INDUCED MINOR EDIT and H -INDUCED TOPOLOGICAL MINOR EDIT are polynomially equivalent to H -INDUCED MINOR and H -INDUCED TOPOLOGICAL MINOR, respectively. We therefore do not consider H -INDUCED MINOR EDIT and H -INDUCED TOPOLOGICAL MINOR EDIT, but will focus on the H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT problems from now on. For these two problems edge deletions are permitted, and this complicates the situation. For example, let $G = K_n$ and $H = K_1$. Then a minimum H -minor sequence of G consists of $n - 1$ vertex deletions, whereas the sequence that consists of $n(n - 1)/2$ edge deletions followed by $n - 1$ vertex deletions is an H -minor sequence of G that has length $n(n - 1)/2 + n - 1$.

Our Results. In Section 2 we pinpoint a close relationship between H -MINOR EDIT and H -INDUCED MINOR, and also between H -TOPOLOGICAL MINOR EDIT and H -INDUCED TOPOLOGICAL MINOR. We use this observation in Section 3.1, where we show both polynomial-time solvable and NP-complete cases for H -

MINOR EDIT and H -TOPOLOGICAL MINOR EDIT; note that the hardness results are in contrast with the aforementioned tractable results for H -MINOR [25] and H -TOPOLOGICAL MINOR [13]. There is currently not much hope in settling the complexity of H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT for all graphs H , due to their strong connection to H -INDUCED MINOR and H -INDUCED TOPOLOGICAL MINOR, the complexity classification of each of which still must be completed. However, in Section 3.2, we are able to show that H -MINOR EDIT is polynomial-time solvable on AT-free graphs, chordal graphs and planar graphs. In Section 3.3 we discuss parameterized complexity aspects, whereas Section 4 contains our conclusions and directions for further research.

Proofs of theorems and lemmas that are marked with the symbol \star have been omitted either completely or partially due to space restrictions.

2 Preliminaries

In this section we state some results from the literature and make some basic observations; we will need these results and observations later on. We only consider undirected finite graphs with no loops and with no multiple edges. We denote the vertex set and edge set of a graph G by V_G and E_G , respectively. If no confusion is possible, we may omit subscripts. We refer the reader to the textbook of Diestel [5] for any undefined graph terminology.

The *disjoint union* of two graphs G and H with $V_G \cap V_H = \emptyset$ is the graph $G + H$ that has vertex set $V_G \cup V_H$ and edge set $E_G \cup E_H$. We let P_n and C_n denote the path and cycle on n vertices, respectively, whereas $K_{1,n}$ is the star on $n + 1$ vertices; note that $K_{1,1} = P_2$ and $K_{1,2} = P_3$. The subgraph of a graph $G = (V, E)$ induced by a subset $S \subseteq V$ is denoted by $G[S]$. A subgraph G' of a graph G is *spanning* if $V_{G'} = V_G$. Let G be a graph that contains a cycle C as a subgraph. If $|V_C| = |V_G|$ then C is a *hamilton cycle*, and G is called *hamiltonian*. An edge $uv \in E_G \setminus E_C$, with C some cycle and with $u, v \in V_C$, is a *chord* of C .

We will frequently make use of the following observation.

Lemma 4. *If (G, k) is a yes-instance of H -MINOR EDIT or H -TOPOLOGICAL MINOR EDIT, for some graph H , then $|V_H| \leq |V_G| \leq |V_H| + k$.*

Proof. Let (G, k) be a yes-instance of H -MINOR EDIT or H -TOPOLOGICAL MINOR EDIT for some graph H . An edge contraction, vertex deletion or vertex dissolution reduces a graph by exactly one vertex, whereas an edge deletion does not change the number of vertices. This has the following two implications. First, no graph operation involved increases the number of vertices of a graph. Hence, $|V_H| \leq |V_G|$. Second, any H -minor sequence of G has length at least $|V_G| - |V_H|$. Hence, $|V_G| - |V_H| \leq k$, or equivalently, $|V_G| \leq |V_H| + k$. \square

We write $\Pi_1 \leq \Pi_2$, for two decision problems Π_1 and Π_2 , to denote that Π_2 generalizes Π_1 . The following observation shows a close relationship between our two editing problems and the corresponding induced containment problems.

Lemma 5. *Let H be a graph. Then the following two statements hold:*

- (i) H -INDUCED MINOR \leq H -MINOR EDIT.
- (ii) H -INDUCED TOPOLOGICAL MINOR \leq H -TOPOLOGICAL MINOR EDIT.

Proof. We start with the proof of (i). Let H be a graph, and let G be an instance of H -INDUCED MINOR. We define $k = |V_G| - |V_H|$. We will show that (G, k) is an equivalent instance of H -MINOR EDIT. If $k < 0$, then G is a no-instance of H -INDUCED MINOR, and by Lemma 4, (G, k) is a no-instance of H -MINOR EDIT. Suppose that $k \geq 0$. We claim that G contains H as an induced minor if and only if (G, k) has an H -minor sequence of length at most k .

First suppose G contains H as an induced minor. Because one edge contraction or one vertex deletion reduces a graph by exactly one vertex, any H -induced minor sequence of G is an H -minor sequence of G that has length $|V_G| - |V_H| = k$.

Now suppose that G has an H -minor sequence S of length at most k . Because $|V_G| - |V_H| = k$, we find that S contains at least k operations that are vertex deletions or edge contractions. Because S has length at most k , this means that S contains no edge deletions. Hence, S is an H -induced minor sequence. We conclude that G contains H as an induced minor.

The proof of (ii) uses the same arguments as the proof of (i); in particular any vertex dissolution in a graph reduces the graph by exactly one vertex. \square

Two disjoint vertex subsets U and W of a graph G are *adjacent* if there exists some vertex in U that is adjacent to some vertex in W . The following alternative definition of being a minor is useful. Let G and H be two graphs. An *H -witness structure* \mathcal{W} is a vertex partition of a subgraph G' of G into $|V_H|$ nonempty sets $W(x)$ called (*H -witness*) *bags*, such that

- (i) each $W(x)$ induces a connected subgraph of G , and
- (ii) for all $x, y \in V_H$ with $x \neq y$, bags $W(x)$ and $W(y)$ are adjacent in G if x and y are adjacent in H .

An *H -witness structure* \mathcal{W} corresponds to at least one H -minor sequence of G . In order to see this, we can first delete all vertices of $V_G \setminus V_{G'}$, then modify all bags $W(x)$ into singletons via edge contractions and finally delete all edges uv with $u \in W(x)$ and $v \in W(y)$ whenever $uv \notin E_H$. The remaining graph is isomorphic to H . We note that G may have more than one H -witness structure.

An *edge subdivision* is the operation that removes an edge uv of a graph and adds a new vertex w adjacent (only) to u and v . This leads to an alternative definition of being a topological minor, namely that a graph G contains a graph H as a topological minor if and only if G contains a subgraph H' that is a *subdivision* of H ; that is, H' can be obtained from H by a sequence of edge subdivisions. A *subdivided star* is a graph obtained from a star after p edge subdivisions for some $p \geq 0$.

Let G be a graph that contains a graph H as a minor. An H -minor sequence S of G is called *nice* if S starts with all its vertex deletions, followed by all its edge contractions and finally by all its edge deletions. It is called *semi-nice* if S starts with all its vertex deletions, followed by all its edge deletions and finally by

all its edge contractions. By replacing edge contractions with vertex dissolutions, we obtain the notions of a *nice* and a *semi-nice* topological minor sequence.

Lemma 6 (★). *Let H be a graph and k an integer. If a graph G has an H -minor sequence of length k , then G has a nice H -minor sequence of length at most k .*

We note that a lemma for topological minors similar to Lemma 6 does not hold. For example, build G as follows: take two disjoint copies of K_n , where $n \geq 5$; subdivide an edge in each copy, to introduce new vertices u and v ; and join the two new vertices u and v . Build H as two disjoint copies of K_n . If we delete the edge (u, v) of G and next perform two vertex dissolutions (of u and v) then we obtain an H -topological minor sequence of length 3 for G . It is not difficult to see that there is no nice H -topological minor sequence for G of length at most 3. However, for topological minor sequences the following holds.

Lemma 7 (★). *Let H be a graph and k an integer. If a graph G has an H -topological minor sequence of length k , then G has a semi-nice H -topological minor sequence S of length at most k , such that the vertices not deleted by the vertex deletions of S induce a subgraph that contains a subdivision of H as a spanning subgraph.*

We note that a lemma for minors similar to Lemma 7 does not hold. The following example, which we will use later on as well, illustrates this.

Example 1. Let $H = C_6$. We take a cycle C_r for some integer $r \geq 7$. Let u be one of its vertices. Add an edge between u and every (non-adjacent) vertex of the cycle except the two vertices at distance two from u . This yields the graph G . Then $(G, r - 5)$ is a yes-instance of H -MINOR EDIT (via a sequence of $r - 6$ edge contractions followed by an edge deletion). It is not difficult to see that every H -minor sequence of length $r - 5$ of G must start with $r - 6$ edge contractions followed by one edge deletion.

Let $K_{k,\ell}$ be the complete bipartite graph with partition classes of size k and ℓ . Fellows et al. [7] showed that for all graphs H , H -INDUCED MINOR is polynomial-time solvable on planar graphs, that is, graphs that contain neither $K_{3,3}$ nor K_5 as a minor. This result has been extended by van 't Hof et al. [18] to any minor-closed graph class that is *nontrivial*, i.e., that does not contain all graphs.

Lemma 8 ([18]). *Let \mathcal{G} be any nontrivial minor-closed graph class. Then, for all graphs H , the H -INDUCED MINOR problem is polynomial-time solvable on \mathcal{G} .*

An *asteroidal triple* in a graph is a set of three mutually non-adjacent vertices such that each two of them are joined by a path that avoids the neighborhood of the third, and *AT-free* graphs are exactly those graphs that contain no such triple. A graph is *chordal* if it contains no induced cycle on four or more vertices. We will also need the following two results.

Lemma 9 ([11]). *For all graphs H , the H -INDUCED MINOR problem can be solved in polynomial time on AT-free graphs.*

Lemma 10 ([2]). *For all graphs H , the H -INDUCED MINOR problem can be solved in polynomial time on chordal graphs.*

3 Complexity Results

In Section 3.1 we consider general input graphs G , whereas in Section 3.2 we consider special classes of input graphs. In Section 3.3 we discuss parameterized complexity aspects.

3.1 General Input Graphs

We first show that the computational complexities of H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT may differ from those of H -MINOR and H -TOPOLOGICAL MINOR, respectively.

Theorem 1. *The following two statements hold:*

- (i) *There is a graph H for which H -MINOR EDIT is NP-complete.*
- (ii) *There is a graph H for which H -TOPOLOGICAL MINOR EDIT is NP-complete.*

Proof. For (i) we take the graph H^* displayed in Figure 1. Then the claim follows from Lemma 2 combined with Lemma 5-(i). For (ii) we take $H = K_5$. Then the claim follows from Lemma 3 combined with Lemma 5-(ii). \square

The remainder of this section is devoted to results for some special classes of target graphs H . We start by considering the case when H is a complete graph; note that Theorem 2-(ii) generalizes Theorem 1-(ii).

Theorem 2. *The following two statements hold:*

- (i) *K_r -MINOR EDIT can be solved in cubic time for all $r \geq 1$.*
- (ii) *K_r -TOPOLOGICAL MINOR EDIT can be solved in polynomial time, if $r \leq 3$, and is NP-complete, if $r \geq 5$.*

Proof. We first prove (i). Let (G, k) be an instance of K_r -MINOR EDIT. If $|V_G| - r < 0$ or $|V_G| - r > k$, then (G, k) is a no-instance of K_r -TOPOLOGICAL MINOR EDIT due to Lemma 4. Suppose that $0 \leq |V_G| - r \leq k$. Because we may remove without loss of generality any edge deletions from a K_r -minor sequence of a graph, we find that (G, k) is a yes-instance of K_r -MINOR EDIT if and only if G contains K_r as a minor. Hence, the result follows after applying Lemma 1.

We now prove (ii). The cases $H = K_1$ and $H = K_2$ are trivial. The case $H = K_3 = C_3$ follows from Theorem 5, which we will prove later. The case $H = K_5$ follows from the proof of Theorem 1-(ii).

Let $r \geq 6$ and assume that K_{r-1} -TOPOLOGICAL MINOR EDIT is NP-complete. Let (G, k) be an instance of K_{r-1} -TOPOLOGICAL MINOR EDIT. From G we construct a graph G' by adding a new vertex v that we make adjacent to all vertices of G . Trivially, if (G, k) is a yes-instance of K_{r-1} -TOPOLOGICAL MINOR EDIT then (G', k) is a yes-instance of K_r -TOPOLOGICAL MINOR EDIT. Conversely,

suppose that (G', k) is a yes-instance of K_r -TOPOLOGICAL MINOR EDIT. Then G' has a K_r -topological minor sequence S of length at most k . We modify S as follows. We remove all edge deletions involving v . We replace the dissolution of any vertex $w \neq v$ that involves v with the deletion of w . If v is not deleted by S , then this yields a K_r -topological minor sequence of G' with length at most k that is a K_{r-1} -topological minor sequence of G with length at most k . If v is deleted by S , then we modify S further by removing this vertex deletion. Because v is adjacent to all vertices of G' other than itself, the resulting sequence is a K_{r+1} -topological minor sequence of G' with length at most $k - 1$. By extending this sequence with the deletion of one of the remaining vertices not equal to v , we obtain a K_r -topological minor sequence of G' with length at most k that is a K_{r-1} -topological minor sequence of G with length at most k . \square

We now consider H -MINOR EDIT for the case in which H is a path or a star.

Theorem 3 (★). P_r -MINOR EDIT and $K_{1,r}$ -MINOR EDIT can both be solved in polynomial time for all $r \geq 1$.

For topological minors we can show a stronger result than Theorem 3.

Theorem 4. Let H be a subdivided star. Then H -TOPOLOGICAL MINOR EDIT is polynomial-time solvable.

Proof. Let H be a subdivided star. Let (G, k) be an-instance of H -TOPOLOGICAL MINOR EDIT with $|V_G| - |V_H| = k'$. We run the algorithm described below.

If $k' < 0$ or $k' > k$, then we return **no**. Otherwise, that is, if $0 \leq k' \leq k$, we proceed as follows. We consider each subset U of $|V_H|$ vertices of G . We check if $G[U]$ contains an H -topological minor sequence of length at most $k - k'$ (note that such a sequence only involves edge deletions). If so, then we return **yes**. Otherwise, after considering all such subsets U , we return **no**.

We now analyze the running time. Let $|V_G| = n$. Then there are at most $n^{|V_H|}$ possible choices of subgraphs G' on $|V_H|$ vertices. This is a polynomial number, because of our assumption that H is fixed. By the same assumption, every such subgraph G' has constant size, and hence can be processed in polynomial time. We conclude that the total running time is polynomial.

We now prove the correctness of our algorithm. If $k' < 0$ or $k' > k$, then (G, k) is a no-instance of H -TOPOLOGICAL MINOR EDIT due to Lemma 4. Suppose that $0 \leq k' \leq k$. We claim that our algorithm returns **yes** if and only if G has an H -topological minor sequence of length at most k .

First suppose that our algorithm returns **yes**. Then G contains H as a topological minor and our algorithm has obtained H from a graph $G[U]$ on $|V_H|$ vertices by at most $k - k'$ operations (which are all edge deletions), whereas the total number of vertex deletions is $|V_G| - |V_H| = k'$. Hence, G has an H -topological minor sequence of length at most $k - k' + k' = k$.

Now suppose that G has an H -topological minor sequence S of length at most k . By Lemma 7, we may assume without loss of generality that S is a semi-nice H -topological minor sequence S of length at most k , such that the

vertices not deleted by the vertex deletions of S induce a subgraph G' that contains a subdivision H' of H as a spanning subgraph. We also assume without loss of generality that the number of vertex deletions in S is maximum over all such H -topological minor sequences of G . Then, because H is a subdivided star, H' must be isomorphic to H (to see this, note that after all vertex deletions of S , we must have a subdivided star, together with possibly some additional edges, and that any subsequent vertex dissolutions can be replaced with vertex deletions so contradicting the maximality of the number of vertex deletions in S). Because $H' \simeq H$ is a spanning subgraph of G' , our algorithm will consider G' at some point. Because the remaining operations in S are at most $k - k'$ edge deletions, they form an H -topological minor sequence of G' that has length at most $k - k'$. This will be detected by our algorithm, which will then return **yes**. This completes the proof of Theorem 4. \square

Note that Theorem 3 can be generalized to be valid for target graphs H that are linear forests (disjoint unions of paths) and Theorem 4 to be valid for target graphs H that are forests, all connected components of which have at most one vertex of degree at least 3, respectively.

We now consider the case when the target graph H is a cycle and show the following result, which holds for topological minors only.

Theorem 5. C_r -TOPOLOGICAL MINOR EDIT can be solved in polynomial time for all $r \geq 3$.

Proof. Let $r \geq 3$. Let (G, k) be an instance of C_r -TOPOLOGICAL MINOR EDIT. We run the following algorithm. Let $k' = |V_G| - r$. If $k' < 0$ or $k' > k$, then we return **no**. Otherwise, we do as follows. We check if G contains C_r as an induced topological minor. If so, then we return **yes**. If not, then we do as follows for each subgraph G' of G with $r \leq |V_{G'}| \leq 2r$. We check if G' contains a C_r -topological minor sequence of length at most $k - (|V_G| - |V_{G'}|)$. If so, then we return **yes**. Otherwise, after having considered all subgraphs G' of G on at most $2r$ vertices, we return **no**.

We first analyze the running time of this algorithm. Checking whether G contains C_r as an induced topological minor is equivalent to checking whether G contains an induced cycle of length at least r ; the latter can be done in polynomial time. Suppose G does not contain C_r as an induced topological minor. Then the algorithm considers at most $|V_G|^{2r}$ subgraphs of G , which is a polynomial number because r is fixed. For the same reason, our algorithm can process every such subgraph G' in constant time.

We are left to prove correctness. If $k' < 0$ or $k' > k$, then (G, k) is a no-instance of C_r -TOPOLOGICAL MINOR EDIT due to Lemma 4. Suppose that $0 \leq k' \leq k$. We claim that our algorithm returns **yes** if and only if G has a C_r -topological minor sequence of length at most k .

First suppose that our algorithm returns **yes**. If G contains C_r as an induced topological minor, then G has a C_r -topological minor sequence of length at most $k' \leq k$. Otherwise, G contains a subgraph G' of at most $2r$ vertices that has a C_r -topological minor sequence of length at most $k - (|V_G| - |V_{G'}|)$. Adding the

$|V_G| - |V_{G'}|$ vertex deletions that yielded G' to this sequence gives us a C_r -topological minor sequence of G that has length at most k .

Now suppose that G has a C_r -topological minor sequence of length at most k . By Lemma 7, we may assume without loss of generality that S is a semi-nice H -topological minor sequence S of length at most k , such that the vertices not deleted by the vertex deletions of S induce a subgraph G' that contains a subdivision H' of C_r as a spanning subgraph; note that $H' = C_s$ for some $s \geq r$. We also assume without loss of generality that the number of vertex deletions in S is maximum over all such H -topological minor sequences of G , and moreover, that C_r is obtained from G' by first deleting all chords and then by dissolving $s - r$ vertices.

If G' has at most $2r$ vertices, then the algorithm would consider G' at some point. Because S is a C_r -topological minor sequence of length at most k , we find that G' has a C_r -topological minor sequence of length at most $k - (|V_G| - |V_{G'}|)$. Hence, our algorithm would detect this and return **yes**.

Now suppose that G' has at least $2r + 1$ vertices. Suppose that C_s has at least one chord e . Because $s = |V_{G'}| \geq 2r + 1$, this means that G' contains a smaller cycle C_t on $t \geq r$ vertices that has exactly $t - 1$ edges in common with C_s . We modify S as follows. We first remove all vertices of G' not on C_t . Then we remove all chords of C_t . Finally, we perform $t - r$ vertex dissolutions on C_t in order to obtain a graph isomorphic to C_r . Hence, the sequence S' obtained in this way is a C_r -topological minor sequence of G as well. By our construction, S' is semi-nice. Moreover, because C_s and C_t have $t - 1$ edges in common, any edge deletion in S' is an edge deletion in S as well. Hence S' has length at most k . However, S' contains at least one more vertex deletion than S , because there exists at least one vertex in G' that is not on C_t . This contradicts the maximality of the number of vertex deletions in S . Hence, C_s has no chords. Then C_s is an induced cycle on at least $s \geq r$ vertices in G' , and consequently, in G . This means that G contains C_r as an induced topological minor. The algorithm checks this and thus returns **yes**. \square

3.2 Input Graphs Restricted to Some Nontrivial Graph Class

Instead of restricting the target graph H to belong to some special graph class, as is done in Section 3.1, we can also restrict the input graph G to some special graph class. In this section we do this for the H -MINOR EDIT problem.

For the H -MINOR EDIT problem, we may use the following lemma that strengthens the relationship between H -MINOR EDIT and H -INDUCED MINOR.

Lemma 11. *Let \mathcal{G} be a graph class and H a graph. If H' -INDUCED MINOR is polynomial-time solvable on \mathcal{G} for each spanning supergraph H' of H , then H -MINOR EDIT is polynomial-time solvable on \mathcal{G} .*

Proof. Let \mathcal{G} be a graph class and H a graph. Suppose that H' -INDUCED MINOR is polynomial-time solvable on \mathcal{G} for each spanning supergraph H' of H . Let $G \in \mathcal{G}$ and $k \in \mathbb{Z}$ form an instance of H -MINOR EDIT. Let $k^* = |V_G| - |V_H|$.

If $k^* < 0$ or $k^* > k$, then we return **no**. Suppose $0 \leq k^* \leq k$. Then, for every spanning supergraph H' of H with at most $k - k^*$ additional edges, we check if G contains H' as an induced minor. As soon as we find that this is the case for some H' we return **yes**. Otherwise, after having considered all spanning supergraphs of H , we return **no**.

The running time of the above algorithm is polynomial for the following two reasons. First, because H is fixed, the number of spanning supergraphs H' of H is a constant. Second, by our assumption, we can solve H' -INDUCED MINOR in polynomial time on \mathcal{G} for each spanning supergraph H' of H .

We now prove that our algorithm is correct. If $k^* < 0$ or $k^* > k$ then (G, k) is a no-instance of H -MINOR EDIT due to Lemma 4. From now on we assume that $0 \leq k^* \leq k$. We claim that our algorithm returns **yes** if and only if G has an H -minor sequence of length at most k .

First suppose that our algorithm returns **yes**. Then there exists a spanning supergraph H' of H with at most $k - k^*$ additional edges, such that H' is an induced minor of G . Let S' be an H' -induced minor sequence of G . Then S' has length exactly $|V_G| - |V_{H'}| = |V_G| - |V_H| = k^*$. We extend S' by deleting the edges in $E_{H'} \setminus E_H$. This yields an H -minor sequence S of G . As $|E_{H'} \setminus E_H| \leq k - k^*$, we find that S has length at most $k^* + k - k^* = k$. Hence, (G, k) is a yes-instance of H -MINOR EDIT.

Now suppose that G has an H -minor sequence S of length at most k . By Lemma 6, we may assume without loss of generality that S is nice, thus all edge deletions in S take place after all its edge contractions and vertex deletions. Let S' be the prefix of S obtained by omitting the edge deletions of S . Then S' is an H' -induced minor sequence of G for some spanning supergraph H' of H . Because every operation in S' is a vertex deletion or edge contraction, S' has length $|V_G| - |V_{H'}| = |V_G| - |V_H| = k^*$. Because S has length at most k , this means that H' can have at most $k - k^*$ more edges than H . Hence, as our algorithm considers all spanning supergraphs of H with at most $k - k^*$ additional edges, it will return **yes**, as desired. This completes the proof of Lemma 11. \square

Combining Lemmas 8–10 with Lemma 11 yields the following result.

Theorem 6. *For all graphs H , the H -MINOR EDIT problem is polynomial-time solvable on*

- (i) *the class of AT-free graphs*
- (ii) *the class of chordal graphs*
- (iii) *any nontrivial minor-closed class of graphs.*

3.3 Parameterized Complexity

A parameterized problem is *fixed-parameter tractable* if an instance (I, p) (where p is the parameter) can be solved in time $f(p) \cdot |V_G|^{O(1)}$ for some function f that only depends on p . Here, the natural parameter is the number of permitted operations k . The following proposition follows immediately from Lemma 4.

Proposition 1. *For all graphs H , the H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT problems are fixed-parameter tractable when parameterized by k .*

We now take $|V_H|$ as the parameter. Theorem 2 shows that in that case H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT are fixed-parameter tractable and para-NP-complete, respectively, when H is a complete graph. The running times of the algorithms given by Theorems 3–5 are bounded by $O(n^{|V_H|})$, where H is a path, subdivided star or cycle, respectively. A natural question would be if we can show fixed-parameter tractability with parameter $|V_H|$ for these cases. However, the following result shows that this is unlikely (the class W[1] is regarded as the parameterized analog to NP).

Proposition 2 (★). *For $H \in \{C_r, P_r, K_{1,r}\}$, the problems H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT are W[1]-hard when parameterized by r .*

Proof. Due to space restrictions, we only consider the cases $H = P_r$ and $H = C_r$.

Let $H = P_r$. Papadimitriou and Yannakakis [24] proved that the problem of testing whether a graph G contains P_r as an induced subgraph is W[1]-hard when parameterized by r (their proof was not done in terms of parameterized complexity theory and was later rediscovered by Haas and Hoffmann [14]). We observe that a graph G contains P_r as an induced subgraph if and only if G contains a P_r -(topological) minor sequence of length at most $|V_G| - r$. Hence, P_r -MINOR EDIT and P_r -TOPOLOGICAL MINOR EDIT are W[1]-hard when parameterized by r .

Let $H = C_r$. It is known that the problem of testing whether a graph G contains C_r as an induced subgraph is W[1]-hard when parameterized by r [14, 24]. The corresponding hardness proofs in [14, 24] immediately imply that the problem of testing whether a graph G contains a cycle C_s with $s \geq r$ as an induced subgraph is W[1]-hard as well, when parameterized by r . We observe that a graph G contains a cycle C_s with $s \geq r$ as an induced subgraph if and only if G contains a C_r -(topological) minor sequence of length at most $|V_G| - r$. Hence, C_r -MINOR EDIT and C_r -TOPOLOGICAL MINOR EDIT are W[1]-hard when parameterized by r . \square

4 Conclusions

The ultimate goal is to complete our partial complexity classifications of H -MINOR EDIT and H -TOPOLOGICAL MINOR EDIT. For this purpose, the following three research questions must be addressed.

1. Is H -MINOR EDIT polynomial-time solvable for all subdivided stars H ?
2. Is C_r -MINOR EDIT polynomial-time solvable for all $r \geq 3$?
3. Is K_4 -TOPOLOGICAL MINOR EDIT polynomial-time solvable?

Answering Question 1 in the affirmative would generalize Theorem 3, in which target graphs H that are paths or stars are considered. Note that generalizing

Theorem 3 to all trees H may be very challenging, because a positive result would solve the aforementioned open problem on H -INDUCED MINOR restricted to trees H , due to Lemma 5-i. The same holds for Question 3: a positive answer to Question 3 would imply membership in P for K_4 -INDUCED TOPOLOGICAL MINOR, the complexity status of which is a notorious open case (see e.g. [21]). As regards Question 2, Example 1 shows that we cannot guess a bounded set of vertices and consider the subgraph that these vertices induce instead of the whole input graph, as was done for C_r -TOPOLOGICAL MINOR EDIT. Hence, new techniques are needed. So far, we only know that the statement is true if $r \leq 4$.

Proposition 3. C_r -MINOR EDIT is polynomial-time solvable if $r \leq 4$.

Proof. If $r = 3$ then $H = C_3 = K_3$ and we apply Theorem 2-(i). Let $r = 4$, and let (G, k) be an instance of C_4 -MINOR EDIT. We run the following algorithm. Let $k' = |V_G| - r$. If $k' < 0$ or $k' > k$, then we return **no** due to Lemma 4. Otherwise, we do as follows. We check if G contains C_4 as an induced minor. If so then we return **yes**. Note that this is equivalent to checking if G contains an induced cycle on at least four vertices, which can be done in polynomial time. If not then G is chordal, and we apply Theorem 6-(ii). \square

Another question is whether we can prove an analog of Theorem 6 for H -TOPOLOGICAL MINOR EDIT. It is known that for all graphs H , the H -INDUCED TOPOLOGICAL MINOR problem is polynomial-time solvable for AT-free graphs [12], chordal graphs [2] and planar graphs [19]. However, as noted in Section 2, we cannot always guarantee the existence of a nice topological minor sequence of sufficiently small length. Hence, our proof technique used to prove Theorem 6 can no longer be applied.

Finally, we can consider other graph containment relations as well. An *edge lift* removes two edges uv and vw that share a common vertex v and adds an edge between the other two vertices u and w involved (should this edge not exist already). A graph G contains a graph H as an *immersion* if G can be modified into H by a sequence of operations consisting of vertex deletions, edge deletions and edge lifts. If edge deletions are not allowed then G contains H as an *induced immersion*. It is known that the corresponding decision problems H -IMMERSION [13] and H -INDUCED IMMERSION [3] polynomial-time solvable for all fixed graphs H (the first problem can even be solved in cubic time [13]). What is the computational complexity of H -IMMERSION EDIT and H -INDUCED IMMERSION EDIT? The main difficulty is that for both problems, we can swap neither edge lifts with vertex deletions nor edge deletions with vertex deletions.

References

1. N. Alon, A. Shapira and B. Sudakov, Additive approximation for edge-deletion problems. In: Proc. FOCS 2005, IEEE Computer Society, 419–428.
2. R. Belmonte, P. A. Golovach, P. Heggernes, P. van 't Hof, M. Kamiński, and D. Paulusma, Detecting fixed patterns in chordal graphs in polynomial time, Algorithmica, to appear.

3. R. Belmonte, P. van 't Hof, and M. Kamiński, Induced Immersions. In: Proc. ISAAC 2012, LNCS 7676 (2012) 299–308.
4. P. Burzyn, F. Bonomo and G. Duran, NP-completeness results for edge modification problems, *Discrete Applied Mathematics* 154 (2006) 1824–1844.
5. R. Diestel, *Graph Theory*, Springer-Verlag, Electronic Edition, 2005.
6. R. G. Downey and M. R. Fellows. Parameterized complexity. Monographs in Computer Science, Springer-Verlag, New York, 1999.
7. M. R. Fellows, J. Kratochvíl, M. Middendorf, and F. Pfeiffer, The complexity of induced minors and related problems, *Algorithmica* 13 (1995) 266–282.
8. J. Fiala, M. Kamiński and D. Paulusma, Detecting induced star-like minors in polynomial time, *Journal of Discrete Algorithms* 17 (2012) 74–85.
9. P.A. Golovach, P. van 't Hof and D. Paulusma, Obtaining planarity by contracting few edges, *Theoretical Computer Science* 476 (2013) 38–46.
10. P.A. Golovach, M. Kamiński, D. Paulusma and D.M. Thilikos, Increasing the minimum degree of a graph by contractions, *Theoretical Computer Science*, to appear.
11. P.A. Golovach, D. Kratsch, D. Paulusma, Detecting induced minors in AT-free graphs, *Theoretical Computer Science*, to appear.
12. P.A. Golovach, D. Paulusma and E.J. van Leeuwen, Induced disjoint paths in AT-free graphs, In: Proc. SWAT 2012, LNCS 7357 (2012) 153–164.
13. M. Grohe, K. Kawarabayashi, D. Marx, and P. Wollan, Finding topological subgraphs is fixed-parameter tractable, In: Proc. STOC 2011, 479–488.
14. R. Haas and M. Hoffmann. Chordless paths through three vertices. *Theoretical Computer Science*, 351 (2006), 360–371.
15. P. Heggernes, P. van 't Hof, B. Lévêque, and C. Paul, Contracting chordal graphs and bipartite graphs to paths and trees, *Discrete Applied Mathematics*, to appear.
16. P. Heggernes, P. van 't Hof, D. Lokshtanov, and C. Paul. Obtaining a bipartite graph by contracting few edges, In: Proc. FSTTCS 2011, 217–228.
17. P. Heggernes, P. van 't Hof, B. Lévêque, D. Lokshtanov, and C. Paul. Contracting graphs to paths and trees, *Algorithmica*, to appear.
18. P. van 't Hof, M. Kamiński, D. Paulusma, S. Szeider and D.M. Thilikos, On graph contractions and induced minors, *Discrete Applied Mathematics* 160 (2012) 799–809.
19. Y. Kobayashi and K. Kawarabayashi. A linear time algorithm for the induced disjoint paths problem in planar graphs. *Journal of Computer and System Sciences* 78 (2012) 670–680.
20. B. Lévêque, D. Y. Lin, F. Maffray, and N. Trotignon, Detecting induced subgraphs, *Discrete Applied Mathematics* 157 (2009) 3540–3551.
21. B. Lévêque, F. Maffray, and N. Trotignon, On graphs with no induced subdivision of K_4 , *Journal of Combinatorial Theory, Series B* 102 (2012) 924–947.
22. J.M. Lewis and M. Yannakakis, The node-deletion problem for hereditary properties is NP-complete, *Journal of Computer and System Sciences* 20 (1980) 219–230.
23. A. Natanzon, R. Shami and R. Sharan, Complexity classification of some edge modification problems, *Discrete Applied Mathematics* 113 (2001) 109–128.
24. C. H. Papadimitriou and M. Yannakakis. On limited nondeterminism and the complexity of the VC-dimension. *Journal of Computer and System Sciences*, 43 (1991), 425–440.
25. N. Robertson and P. D. Seymour, Graph minors. XIII. The disjoint paths problem. *Journal of Combinatorial Theory, Series B* 63 (1995) 65–110.